

On the Bound States of Matrix Strings

Vatche Sahakian
Enrico Fermi Institute, Department of Physics
University of Chicago
5640 S. Ellis Ave., Chicago, IL 60637, USA
`isaak@maxwell.uchicago.edu`

Abstract

We investigate excitations in Matrix Theory on T^2 corresponding to bound states of strings. We demonstrate the Dirichlet aspect of R-R charged vacua through a non-trivial connection between the $U(1)$ and $SU(n)$ sectors of the matrix SYM.

1 Introduction

During the past few years, a web of perturbative and non-perturbative connections has been cast between the different string theories of a decade ago[1]; the common theme is that a limiting regime of a given theory is described by another. Probing the non-perturbative regime of the IIA string theory, with 11 dimensional supergravity as a description of its low energy dynamics, M-theory[2] stands at a crucial junction in the web of string dualities, with the need of a proper description of its degrees of freedom. The BFSS conjecture[3] proposes that the full dynamics of M theory in the infinite momentum frame is captured by the matrix SYM describing IIA D0 branes. Given the already established connections between M and other theories in various settings, this proposal faces a multitude of consistency tests. And many it has already, and remarkably, passed[4]. Of relevance to the current discussion, it is expected, on the basis of proposed dualities, to account for the bound states of IIB strings[5][6] in this Matrix Theory. It was argued in [7][8] that, Matrix Theory on T^1 , a 1 + 1 SYM, metamorphoses, in the strong YM coupling regime, into the CFT of the IIA string in the infinite momentum frame; this entailed an elegant mechanism of “screwing” string-like excitations into matrices. The first quantized IIB fundamental string was identified along a similar mechanism in Matrix Theory on T^2 in [9]; it was proposed that the $SL(2, Z)$ S-duality of the IIB theory can then be used to realize the spectrum of the IIB bound strings. The purpose of this paper is to investigate this latter point in greater detail.

We will study the energetics of excitations in Matrix Theory on the torus. Storing NS-NS and R-R charges in the $U(1)$ sector, we will propose to wind the dual gauge scalar of the theory on a cycle determined by the charge content of this vacuum. The energetics of excitations in the $SU(n)$ will be seen to be affected by the $U(1)$ configuration in a non-trivial pattern; quanta parallel to the $U(1)$ electric field will become light if the latter corresponds to an R-R charged configuration; these light excitations will be indicative of massive excitations of fundamental strings attached to the background, a signature of the Dirichlet aspect of these vacua[10][11]. The heavy excitations in these scenarios will be perpendicular to the electric field and will be identified with the non-BPS excitations of IIB bound strings.

Throughout, we will be sloppy in handling factors of 2π ; the reader may set $2\pi \rightarrow 1$ to avoid grief.

2 Matrix Theory on T^d

Consider Matrix Theory as M theory in the infinite momentum frame described by the quantum mechanics Lagrangian[3]

$$L = \frac{1}{2R_{11}} tr \dot{X}_i^2 - \frac{R_{11}}{2} \frac{1}{l_P^6} tr [X_i, X_j]^2 + fermions , \quad (1)$$

where the 9 X ’s are hermitian $N \times N$ matrices and the quantum commutator is given by

$$[X_{ab}^i, X_{cd}^j] = iR_{11}\delta_{ij}\delta_{ac}\delta_{bd} . \quad (2)$$

The dimensional reduction of this theory on a transverse torus T^d can be obtained by realizing a subset of the $U(N)$ gauge symmetry as the compactification of d scalars, yielding a $d+1$ SYM[3][12]. After rescaling fields and coordinates, the form we will use is

$$\mathcal{L} = \frac{1}{2} \left(\frac{l_P^2}{R_{11}} \right)^{3-d} tr \left\{ -\frac{F_{\mu\nu}^2}{(R_\mu R_\nu)^2} + \frac{(D_\mu X_i)^2}{R_\mu^2} - G^2 [X_i, X_j]^2 + fermions \right\} , \quad (3)$$

with

$$F_{\mu\nu} \equiv R_\nu \partial_\mu A_\nu - R_\mu \partial_\nu A_\mu - i R_\mu R_\nu G[A_\mu, A_\nu] \quad (4)$$

$$D_\mu \equiv \partial_\mu - i R_\mu G[A_\mu, \cdot] , \quad (5)$$

where $\mu, \nu = 0, \dots, d$, and $i, j = d+1, \dots, 9$; note that repeated indices in some of these equations are not summed over. The dimensionless coupling G is

$$G^2 \equiv \frac{l_P^d}{\prod_\mu R_\mu} . \quad (6)$$

The quantum commutator for the gauge or scalar fields becomes

$$[X_{ab}^i(\sigma), \dot{X}_{cd}^j(\sigma')] = \left(\frac{l_P^2}{R_{11}} \right)^{d-3} \prod_\mu \frac{\delta(\sigma^\mu - \sigma'^\mu)}{R_\mu} \delta_{ij} \delta_{ac} \delta_{bd} . \quad (7)$$

This YM is on the torus parametrized by σ^μ for $\mu = 1, \dots, d$, with periods

$$\Sigma^\mu = \frac{l_P^3}{R_{11} R_\mu^2} , \quad (8)$$

and the measure of the action integral is

$$d\tau \prod_\mu d\sigma^\mu R_\mu . \quad (9)$$

Some of the equations have time $\tau \equiv \sigma^0$ rescaled by an $R^0 \sim 1$; no significance beyond notational convenience will be accorded to this scale. The fermions are ignored throughout; supersymmetry can be invoked to reintroduce their contributions to the final results.

We will also need some of the charges appearing in the SUSY algebra[8]; in our notation, these are

$$\oint tr F_{0\mu} = w q_\mu \frac{R_{11} R_0}{G R_\mu^2} \quad (10)$$

$$\oint tr F_{\mu\nu} = \frac{w_{\mu\nu}}{G} , \quad (11)$$

where w , q_μ and $w_{\mu\nu}$ are integers by momentum and/or Dirac quantization; $w q_\mu$ is momentum in the appropriate direction, or Kaluza charge equivalently, while $w_{\mu\nu}$ counts the winding of the 11D membrane on corresponding 2-cycles. We have also factored out a common integer w in the momenta; in the case $d = 2$, this implies that q_1 and q_2 are relatively prime.

We will consider this system for $d = 2$, i.e. SYM 2 + 1 as M-theory on a transverse torus. Taking $R_{11} \rightarrow \infty$ as part of the infinite momentum frame prescription, the three length scales of this theory are R_1 , R_2 and l_P . We focus on the strong coupling regime of the YM

$$G^2 \gg 1 \Rightarrow R_1 R_2 \ll l_P^2 . \quad (12)$$

Energy considerations then justify dropping excitations in non-commuting matrices; we then go to the diagonal gauge where we have the residual gauge freedom corresponding to the permutations of matrix eigenvalues[7][8]. Furthermore, we can now dualize the field strength [9] into the derivative of a scalar

$*F = d\phi$; we will do this in the coordinates $R_\mu \sigma_\mu$; rescaling the scalar as well for convenience, we define the duality map

$$\dot{\phi} = \frac{GR_0}{R_1 R_2} F_{12} \quad (13)$$

$$\partial_1 \phi = \frac{GR_1}{R_0 R_2} F_{02} \quad (14)$$

$$\partial_2 \phi = -\frac{GR_2}{R_0 R_1} F_{01} . \quad (15)$$

In the regime we are considering, the Hamiltonian then takes the form

$$H = \frac{1}{2R_{11}} \int d^2 \sigma \text{tr} \left(R_1^2 (\partial_2 \phi)^2 + R_2^2 (\partial_1 \phi)^2 + R_1^2 R_2^2 \dot{\phi}^2 + R_1^2 (\partial_2 X^i)^2 + R_2^2 (\partial_1 X^i)^2 + R_1^2 R_2^2 \dot{X}^i{}^2 \right) , \quad (16)$$

where $i = 3, \dots, 9$, and the X 's have been rescaled $X \rightarrow X/G$; the quantum commutator for either the ϕ or the X 's is

$$[\phi_{aa}(\sigma), \dot{\phi}_{bb}(\sigma')] = iR_{11} \frac{\delta^2(\sigma - \sigma')}{R_1^2 R_2^2} \delta_{ab} . \quad (17)$$

3 Exciting matrices

We want to look for excitations in the Hamiltonian of equation (16) corresponding to the bound states of fundamental and Dirichlet strings, predicted to be in the spectrum through the duality between M-theory on T^2 and the IIB theory on the circle[13]. We focus on a block diagonal matrix configuration of size n sitting in the larger $N \times N$ matrices; the size of this block is expected to be identified with the longitudinal momentum of excitations it will carry, and is to be taken to infinity.

We excite zero modes in the $U(1)$ gauge fields according to equations (10) and (11); we then have an electric fields along the (q_1, q_2) cycle and a 2-form magnetic field on the torus. It easy to show that excitations in the $U(1)$ apart from these zero modes contribute mass $\propto n \rightarrow \infty$ ¹. Throwing away such non-zero modes in the $U(1)$, we find that the scalar is given by

$$\phi^{U(1)} = \frac{w_{12}}{n} \frac{R_{11}^2 R_1 R_2 R_0}{l_P^6} \tau + \frac{w}{n} q_2 \frac{R_{11}^2 R_1}{l_P^3 R_2} \sigma^1 - \frac{w}{n} q_1 \frac{R_{11}^2 R_2}{l_P^3 R_1} \sigma^2 . \quad (18)$$

Along with zero modes in the X 's, this gives the $U(1)$ energy

$$H^{U(1)} = \frac{R_{11}}{2n} \left(w^2 \left(\frac{q_1^2}{R_1^2} + \frac{q_2^2}{R_2^2} \right) + \left(\frac{w_{12} R_1 R_2}{l_P^3} \right)^2 + p_i^2 \right) . \quad (19)$$

The charges stored in the $U(1)$ will eventually be interpreted as R-R or NS-NS charges in the IIB theory. Given the connection between R-R charges and the Dirichlet aspect of objects carrying such charges, we expect a non-trivial mechanism through which the $U(1)$ sector of the YM affects the nature of the excitations in the $SU(n)$ sector; this despite the fact that, superficially, the two sectors are decoupled. We will demonstrate this connection by looking at the theory on a transformed torus.

¹This can be seen as follows; the non-zero modes in the $U(1)$ will have period Σ ; two derivatives in the energy and the trace of the unit matrix contribute collectively a factor of n to the Hamiltonian; with the longitudinal momentum given by $p_+ = n/R_{11}$, this corresponds to a mass $\propto n$, which is out of the spectrum in the infinite momentum frame.

We apply the transformation to new coordinates $(\tilde{\sigma}^1, \tilde{\sigma}^2)$ that aligns the one-form electric field along $\tilde{\sigma}^1$; this transformation then must take the form

$$\begin{pmatrix} \tilde{\sigma}^1 \\ \tilde{\sigma}^2 \end{pmatrix} = \begin{pmatrix} q_1 & q_2 \\ c & -d \end{pmatrix} \begin{pmatrix} \sigma^1 \\ \sigma_2 \end{pmatrix}, \quad (20)$$

where we restrict c and d by orthogonality and area preservation

$$c = -\frac{q_2 R_1^2}{\Delta} \quad (21)$$

$$d = -\frac{q_1 R_2^2}{\Delta} \quad (22)$$

$$\Delta \equiv q_1^2 R_2^2 + q_2^2 R_1^2. \quad (23)$$

The $SU(n)$ part of the Hamiltonian of equation (16) becomes

$$H^{SU(n)} = \frac{R_1 R_2}{2R_{11}} \int d^2 \tilde{\sigma} \text{tr} \left(\frac{\Delta}{R_1 R_2} (\tilde{\partial}_1 \phi)^2 + \frac{R_1 R_2}{\Delta} (\tilde{\partial}_2 \phi)^2 + R_1 R_2 \dot{\phi}^2 + \text{similar X terms} \right). \quad (24)$$

In the new variables, the scalar field looks like

$$\phi^{U(1)} = \frac{w_{12}}{n} \left(\frac{R_{11}}{l_P^3} \right)^2 R_1 R_2 \tau - \frac{w}{n} \frac{R_{11}^2 \Delta}{l_P^3 R_1 R_2} \tilde{\sigma}^2. \quad (25)$$

Given that this zero mode structure depends on the coordinate $\tilde{\sigma}^2$, the natural candidate for string-like excitations away from this configuration are modes of the same σ dependence, i.e. perpendicular to the direction of the electric field

$$\phi = \phi^{U(1)}(\tilde{\sigma}^2) + \phi^{SU(n)}(\tilde{\sigma}^2). \quad (26)$$

To find the periods of the new torus, we will again make use of the $U(1)$ sector. The scalar being compact with radius R , we wind $\tilde{\sigma}^2$ on R using equation (25)²

$$\frac{R_{11}^2 \Delta}{l_P^3 R_1 R_2} \tilde{\Sigma}^2 = R. \quad (27)$$

From the quantum commutator (17), we also have

$$R = \frac{R_{11}}{R_1 R_2}, \quad (28)$$

and area preservation implies

$$\tilde{\Sigma}_1 \tilde{\Sigma}_2 = \Sigma_1 \Sigma_2. \quad (29)$$

Putting things together, we find

$$\tilde{\Sigma}_1 = \frac{l_P^3 \Delta}{R_{11} R_1^2 R_2^2} \quad (30)$$

²We note that the transformation aligns the new axis along integer valued vectors on the torus, and therefore yields to new periodic coordinates; this point was brought to my attention by M. Li. We also note that the momenta stored in the $U(1)$ sector are quantized on the original torus, as seen in equation (18); in this respect, the transformation at hand is used as a tool to make manifest the energetics of the excitations.




Figure 1: The YM torus in the regime $R_2 \ll R_1$ (see equation (8) for this identification); the solid arrow is the direction of the $U(1)$ electric field in three different scenarios, from left to right corresponding to $q_2 = 0$, both charges nonzero, and $q_1 = 0$; the dotted arrow is the direction of light excitations in each case.

$$\tilde{\Sigma}_2 = \frac{l_P^3}{R_{11}\Delta} . \quad (31)$$

To analyze the energetics of the $SU(n)$ Hamiltonian, we rescale the coordinates $(\tilde{\sigma}^1, \tilde{\sigma}^2) = (\tilde{\Sigma}^1 \tilde{\sigma}'^1, \tilde{\Sigma}^2 \tilde{\sigma}'^2)$ to new variables of period 1 (or 2π). The $SU(n)$ Hamiltonian then takes the form

$$H^{SU(n)} = \frac{1}{2} \int d^2 \tilde{\sigma}' \text{tr} \left(R_1 R_2 \left(\frac{R_1 R_2}{\Delta} \right) \left(\frac{R_{11}}{l_P^3} \right)^2 (\tilde{\sigma}'_1 \phi)^2 \right. \\ \left. + R_1 R_2 \left(\frac{\Delta}{R_1 R_2} \right) \left(\frac{R_{11}}{l_P^3} \right)^2 (\tilde{\sigma}'_2 \phi)^2 + \dot{\phi}^2 + \dots \right) , \quad (32)$$

where we have rescaled the fields again so as to get the canonical form of the commutator for the ϕ , i.e. schematically $[\phi, \dot{\phi}] = i\delta^2$.

Let us first consider the case where $q_1 \neq 0$ and $q_2 \neq 0$. We then note that the following strict inequality holds

$$\frac{\Delta}{R_1 R_2} > 1 \Rightarrow \tilde{\Sigma}^1 > \tilde{\Sigma}^2 \quad (33)$$

and can be pushed to the extreme by either $R_2 \ll R_1$ or $R_1 \ll R_2$. The electric field is parallel to $\tilde{\sigma}^1$; looking at the relative relations of the σ derivatives in equation (32), we notice that the string-like $SU(n)$ excitations parallel to the $U(1)$ electric field are energetically favored over the more natural candidates perpendicular to it, in *both* regimes $R_2 \ll R_1$ and $R_1 \ll R_2$; we have the non-trivial statement that, in the skewed torus limit, the identification of the direction of the light $SU(n)$ excitations depends on the direction of the $U(1)$ electric field, independent of our choice of the small radius; the content of the $U(1)$ vacuum is affecting the nature of the excitations in the $SU(n)$ due to the existence of a scalar in the theory winding on a circle as in equation (27). Extending this analysis to the zero charge cases as well, the different possible scenarios are illustrated in Figure 1. We note the pattern that, whenever the electric field, indicated by the solid arrow, projects a component along the long side of the torus (i.e. the direction of the small radius), the light excitations flip parallel to it, whereas the natural string-like excitations throughout are expected to be perpendicular to the electric field, as seen from equation (26). We will demonstrate that this structure translates into the link between R-R charge and the Dirichlet aspect of R-R charged vacua.

For the purpose of substantiating the meaning of both the light and heavy excitations in the $SU(n)$ sector, we would like to consider exciting each of the two cycles independently and writing down the corresponding energy spectra. For the $U(1)$ sector, the zero modes are determined and we have the boundary condition

$$\phi^{U(1)}(\tilde{\sigma}^2 + \tilde{\Sigma}^2) = \phi^{U(1)}(\tilde{\sigma}^2) + R \frac{w}{n} \quad (34)$$

which was used in equation (27) already. Next, look at excitations in the $SU(n)$ with Motl-like twisted boundary conditions [7] along either direction $\tilde{\sigma}^1$ or $\tilde{\sigma}^2$. It can be shown that the most general physically relevant boundary condition in the $SU(n)$ sector is

$$\phi^{SU(n)}(\sigma + \Sigma) = g \phi^{SU(n)}(\sigma) g^{-1} , \quad (35)$$

where σ is either $\tilde{\sigma}^1$ or $\tilde{\sigma}^2$ and g is the shift operator in Z_n ³; after rescaling the time appropriately, we then easily find the form⁴

$$\phi^{SU(n)} = i \frac{\alpha_m}{m} \omega^m \exp - \frac{2\pi m i}{\Sigma n} (\tau - \sigma) + i \frac{\tilde{\alpha}_m}{m} \omega^{-m} \exp - \frac{2\pi m i}{\Sigma n} (\tau + \sigma) \quad (37)$$

with $\omega = \text{diag}(e^{2\pi i/j})$. Similar boundary conditions and solutions follow for the other X polarizations. The $SU(n)$ energies for excitations along $\tilde{\sigma}^1$ and $\tilde{\sigma}^2$ are easily read off the Hamiltonian (32)

$$H_{(1)}^{SU(n)} = \frac{R_{11}}{2n} 4\pi \frac{\sqrt{R_1 R_2}}{l_P^3} \left(q_1^2 \frac{R_2}{R_1} + q_2^2 \frac{R_1}{R_2} \right)^{-1/2} \hat{N} \quad (38)$$

$$H_{(2)}^{SU(n)} = \frac{R_{11}}{2n} 4\pi \frac{\sqrt{R_1 R_2}}{l_P^3} \left(q_1^2 \frac{R_2}{R_1} + q_2^2 \frac{R_1}{R_2} \right)^{1/2} \hat{N} . \quad (39)$$

The $\hat{N}$ operator is the usual right and left number operator which may now be thought to include contributions from fermionic excitations.

Consider the regime

$$R_2 \ll R_1 \Rightarrow R_2 \ll l_P . \quad (40)$$

From the point of view of M-theory and its known connections with IIA and IIB string theories, excitations in this regime are expected to be described by a perturbative IIA theory with dilaton vev $e^{\phi_A} = g_A$ and string length scale l_S given by

$$R_2 = g_A l_S \quad (41)$$

$$l_P^3 = l_S^3 g_A \quad (42)$$

³Elements of S_N split into cycles in Z_n such that, by construction, we have focused on a block diagonal matrix mapping into one of these cycles; see [8] for details on this construction.

⁴It is mildly interesting to note that the structure of sewing the strings into these matrices can be shown to forbid zero modes in the $SU(n)$. Generally, the residual gauge freedom in the diagonal gauge consists of matrices of the form $gd(\sigma)$, where $g \in S_N$ and is independent of σ by continuity, while $d(\sigma)$ is a diagonal matrix contributing only through the derivative term in the gauge transformation. The most general boundary condition on the scalar is then

$$\phi^{SU(n)} = g \phi^{SU(n)} g^{-1} + D , \quad (36)$$

where D is a diagonal traceless matrix independent of σ and g is the shift operator in Z_n as argued in the previous footnote. Writing $\phi^{SU(n)} = M_1 + M_2 \sigma + M_3 \tau + \phi_{nonzero}^{SU(n)}$, we can easily show that $M_2 = M_3 = 0$, while M_1 can be gauged away. We are then left with the most general boundary condition of equation (35) on the non-zero modes.

or by a perturbative IIB theory with dilaton vev $e^{\phi_B} = g_B$

$$g_B = R_2/R_1 . \quad (43)$$

The IIB fundamental string tension is then given in terms of M theory parameters by

$$T_F = \frac{R_2}{l_P^3} , \quad (44)$$

while the IIB Dirichlet string tension is

$$T_D = \frac{R_1}{l_P^3} \gg T_F . \quad (45)$$

Furthermore, q_1 gets interpreted as NS-NS charge, while q_2 counts R-R charge. Looking at Figure 1, we note that the configurations with light excitations parallel to the electric field correspond to scenarios carrying non-zero R-R charge.

Consider first the case of $q_1 = 1, q_2 = 0$. The $SU(n)$ sector is excited energetically favorably in the σ^2 direction, perpendicular to the electric field, as shown in Figure 1. Given that $H^{U(1)} \sim \mathcal{O}(T_F)$, the skewed torus limit singles out the light string-like excitations to $\mathcal{O}(T_F)$ in the $SU(n)$ as well. The excitations off the $U(1)$ vacuum are 1+1 dimensional and the spectrum of the Hamiltonian $H = H^{U(1)} + H^{SU(n)}$ is that of the IIB fundamental string[9]

$$H = \frac{R_{11}}{2n} \left((2\pi R_B w T_F)^2 + \left(\frac{w_{12}}{R_B} \right)^2 + p_i^2 + 4\pi T_F \hat{N} \right) , \quad (46)$$

where we have identified the decompactifying IIB radius

$$R_B = \frac{l_P^3}{R_1 R_2} \gg l_P . \quad (47)$$

As soon as we put a single unit of R-R charge in the $U(1)$, i.e. the electric field projects onto the long side in Figure 1, the situation changes in two ways; first, $H^{U(1)} \sim \mathcal{O}(T_D)$; second, the string-like light excitations are now along a parallel direction to the electric field and are of $\mathcal{O}(T_F)$. Considering the tower of winding modes in the $U(1)$, we are not justified to ignore the heavy excitations along $\tilde{\sigma}^2$ from the spectrum. The heavy excitations present the spectrum of the (q_1, q_2) string[6]

$$H = \frac{R_{11}}{2n} \left(p_i^2 + M_{(q_1, q_2)}^2 \right) , \quad (48)$$

where $i = 3, \dots, 9$, and

$$M_{(q_1, q_2)}^2 = (2\pi R_B w T_{(q_1, q_2)})^2 + \left(\frac{w_{12}}{R_B} \right)^2 + 4\pi T_{(q_1, q_2)} \hat{N} . \quad (49)$$

Off each level in this tower, we have a light spectrum

$$H_{(1)}^{SU(n)} \rightarrow \frac{R_{11}}{2n} 4\pi \frac{T_F}{q_2} \hat{N} . \quad (50)$$

In general, it may be argued that a complete description of the system requires the full dynamics of the membrane theory with correspondingly more complicated boundary conditions on the torus. Nevertheless,

we claim that this picture is enough to allow us to summarize the conclusions of the analysis as follows. Given a $U(1)$ electric field on the torus, we obtain the spectrum of the (q_1, q_2) bound strings by exciting the direction perpendicular to the field in the $SU(n)$; configurations carrying zero R-R charge have a tower of heavy excitations parallel to the electric field pushed out of the spectrum to $\mathcal{O}(T_F)$, leaving the spectrum of the fundamental IIB string. Configurations carrying R-R charge, through the non-trivial mechanism outlined above, have the bound string spectrum additionally swamped by light excitations parallel to the electric field. We propose that these light excitations correspond to massive excitations of fundamental strings attached to bound state of strings; even a single unit of R-R charge in the $(0, 1)$ D-string generates this light spectrum; this is the signature of the Dirichlet aspect of the underlying $U(1)$ vacuum. The fact that these light excitations are of different σ dependence than the ones in the $U(1)$ sector indicates further the different role played by these light strings.

4 Conclusions

We have demonstrated, using energy arguments, the mechanism through which excitations in Matrix Theory, essentially excitations on a membrane, distribute themselves into the spectrum of bound IIB strings and the spectrum of fundamental strings attached to D strings. We have also translated the connection between R-R charge and the Dirichlet aspect of the D-strings into the language of the matrix YM, that of a non-trivial connection between the $U(1)$ and the $SU(n)$ sectors. This feature arose from the mechanism of winding a $U(1)$ scalar onto an oblique cycle, given the charge content of the configuration; consequently, the periods of the torus seen by the $SU(n)$ excitations contained the information about the winding $U(1)$ scalar and the charges it carries. Finally, we note that, whereas in [14] the BPS condition picked out the string-like excitations on a cycle of the membrane to be identified with the short spectrum of the bound strings, in the Matrix Theory the energetics of the excitations naturally splits the dynamics on the membrane into two sets of string-like spectra, one of which is mapped onto the full non-BPS spectrum of the bound strings.

To further substantiate this proposal, a picture tracing the geometry and some of the dynamics of the massive excitations of the strings attached to the state of bound strings is desirable; this may possibly explain the exact form of the light excitations we have found.

5 Acknowledgments

I would like to thank E. Martinec for suggesting the problem and for particularly illuminating discussions at various stages of the analysis. I would also like to acknowledge M. Li for useful suggestions.

References

- [1] Lectures on Superstring and M Theory Dualities, John H. Schwarz, hep-th/9607201
- [2] String Theory Dynamics In Various Dimensions, Edward Witten, hep-th/9503124
- [3] M Theory As A Matrix Model: A Conjecture, T. Banks, W. Fischler, S.H. Shenker, L. Susskind, hep-th/9610043
- [4] The State of Matrix Theory, Tom Banks, hep-th/9706168

- [5] Bound States Of Strings And p -Branes, Edward Witten, hep-th/9510135
- [6] An $SL(2, Z)$ Multiplet of Type IIB Superstrings, John H. Schwartz, hep-th/9508143
- [7] Proposals on nonperturbative superstring interactions, Lubos Motl, hep-th/9701025
- [8] Matrix String Theory, R. Dijkgraaf, E. Verlinde, H. Verlinde, hep-th/9703030
- [9] Strings from Matrices, Tom Banks, Nathan Seiberg, hep-th/9702187
- [10] Dirichlet-Branes and Ramond-Ramond Charges, Joseph Polchinski, hep-th/9510017
- [11] TASI Lectures on D-Branes, Joseph Polchinski, hep-th/9611050
- [12] D-brane field theory on compact spaces, Washington Taylor, hep-th/9611042
- [13] Superstring Dualities, John H. Schwarz, hep-th/9509148
- [14] Waves, boosted branes and BPS states in M-theory, J.G. Russo, A.A. Tseytlin, hep-th/9611047